

Comment on ‘On the next-to-leading order gravitational spin(1)-spin(2) dynamics’ by J. Steinhoff et al.

Rafael A. Porto¹ and Ira Z. Rothstein²

¹*Physics Department, University of California, Santa Barbara
Santa Barbara, 93106 USA*

²*Carnegie Mellon University Dept. of Physics,
Pittsburgh PA 15213, USA*

In this comment we explain the discrepancy found between the results in arXiv:0712.1716v1 for the 3PN spin-spin potential and those previously derived in gr-qc/0604099. We point out that to compare one must include sub-leading lower order spin-orbit effects which contribute to the spin-spin potential once one transforms to the PN frame. When these effects are included the results in arXiv:0712.1716v1 do indeed reproduce those found in gr-qc/0604099.

In [1] we applied recently developed Effective Field Theory (EFT) techniques [2, 4] to compute the next to leading order (NLO) potential due to spin(1)-spin(2) interactions in the Newton-Wigner (NW) spin supplementarity condition (SSC). In [5] the NLO spin(1)-spin(2) was also calculated. Given our claim that, up to 4PN order the equations of motion in the spin(1)-spin(2) sector can be obtained via the traditional Hamiltonian approach [1, 6], the authors then searched for the canonical transformation necessary to go between the results, finding that no such transformation exists. However, in order to compare one must also consider spin-orbit effects, including a subleading one which becomes a spin(1)spin(2)-orbit term once written in terms of the coordinate velocity in the post-Newtonian one. In fact, once this new term is added, the result in [5] do indeed reproduce the potential calculated in [1].

The spin-orbit 1.5PN potential is given by [4, 6]

$$V_{1.5PN}^{so} = \frac{G_N m_2}{r^2} n^j \left(S_1^{j0} + S_1^{jk} (v_1^k - 2v_2^k) \right) + 1 \leftrightarrow 2, \quad (1)$$

and therefore, it depends on S^{j0} . As we emphasized before [6], in our formalism the spin variable lives in a locally flat frame, where $S^{ab} = S^{\mu\nu} e_\mu^a e_\nu^b$ and e_μ^a is a local tetrad field such that $e_\mu^a e_\nu^b = g_{\mu\nu}$ [6]. The spin algebra is thus given by the standard $SO(1,3)$, and the spin gravity interaction takes the form, $-\frac{1}{2} \omega_\mu^{ab} S_{ab}$ with ω_μ^{ab} the Ricci rotation coefficients. Expanding around flat space we have $e_\mu^b = \delta_\mu^b + \frac{h_{\nu\mu}}{2} \delta_\nu^b + \dots$, which allows us to obtain the Feynman rules with which the potential is calculated [1, 4, 6].

On the other hand, the NW SSC [7] written in terms of the local spin and the coordinate velocity, $v^j \equiv \frac{dx^j}{dt}$, then reads

$$S_1^{i0} = \frac{1}{2} S_1^{ij} v_1^j + \frac{1}{2} S_1^{ij} e_0^j(\vec{x}_1) + \dots = \frac{1}{2} (\vec{v}_1 \times \vec{S}_1)^i + \frac{G_N}{2r^2} \left((\vec{n} \times \vec{S}_2) \times \vec{S}_1 \right)^i + \dots, \quad (2)$$

where we have $e_0^j = H_0^j/2 = \frac{G_N}{r^2} (\vec{n} \times \vec{S}_2)^j + \dots$, which follows from the one point function, $\langle H_0^j \rangle$, using the leading order spin Feynman rule $\frac{i}{2} h_{0i,j} S^{ij}$ [1, 4], or by simple inspection of the Kerr metric in harmonic gauge. A similar term follows for particle 2.

In the EFT approach [1, 2, 4, 6] each vertex scales with different power of spin, and velocity, and therefore spin(1)-spin(2) interactions are those for which there is a spin tensor on each vertex. On the other hand spin-orbit terms are those for which the spin of one of the bodies interacts with the motion of the companion. As we can see from (1) and (2), to obtain the full $S_1 S_2$ Hamiltonian up to 3PN we need to include a spin(1)spin(2)-orbit term given by

$$\frac{G_N^2}{2r^2} \left(m_2 n^i S_1^{ij} e_0^j(\vec{x}_1) - m_1 n^i S_2^{ij} e_0^j(\vec{x}_2) \right) = \frac{G_N^2 M}{2r^4} \left((\vec{S}_1 \times \vec{n}) \cdot (\vec{n} \times \vec{S}_2) \right) = \frac{G_N^2 M}{2r^4} \left(\vec{S}_1 \cdot \vec{n} \vec{S}_2 \cdot \vec{n} - \vec{S}_1 \cdot \vec{S}_2 \right), \quad (3)$$

where $M = m_1 + m_2$,

and we thus get

$$\begin{aligned}
V^{s1s2} = & -\frac{G_N}{2r^3} \left[\vec{S}_1 \cdot \vec{S}_2 \left(\frac{3}{2} \vec{v}_1 \cdot \vec{v}_2 - 3\vec{v}_1 \cdot \vec{n} \vec{v}_2 \cdot \vec{n} - (\vec{v}_1^2 + \vec{v}_2^2) \right) - \vec{S}_1 \cdot \vec{v}_1 \vec{S}_2 \cdot \vec{v}_2 - \frac{3}{2} \vec{S}_1 \cdot \vec{v}_2 \vec{S}_2 \cdot \vec{v}_1 + \vec{S}_1 \cdot \vec{v}_2 \vec{S}_2 \cdot \vec{v}_2 \right. \\
& + \vec{S}_2 \cdot \vec{v}_1 \vec{S}_1 \cdot \vec{v}_1 + 3\vec{S}_1 \cdot \vec{n} \vec{S}_2 \cdot \vec{n} (\vec{v}_1 \cdot \vec{v}_2 + 5\vec{v}_1 \cdot \vec{n} \vec{v}_2 \cdot \vec{n}) - 3\vec{S}_1 \cdot \vec{v}_1 \vec{S}_2 \cdot \vec{n} \vec{v}_2 \cdot \vec{n} - 3\vec{S}_2 \cdot \vec{v}_2 \vec{S}_1 \cdot \vec{n} \vec{v}_1 \cdot \vec{n} \\
& + 3(\vec{v}_2 \times \vec{S}_1) \cdot \vec{n} (\vec{v}_2 \times \vec{S}_2) \cdot \vec{n} + 3(\vec{v}_1 \times \vec{S}_1) \cdot \vec{n} (\vec{v}_1 \times \vec{S}_2) \cdot \vec{n} - \frac{3}{2} (\vec{v}_1 \times \vec{S}_1) \cdot \vec{n} (\vec{v}_2 \times \vec{S}_2) \cdot \vec{n} \\
& \left. - 6(\vec{v}_1 \times \vec{S}_2) \cdot \vec{n} (\vec{v}_2 \times \vec{S}_1) \cdot \vec{n} \right] + \frac{G_N^2 M}{2r^4} \left(5\vec{S}_1 \cdot \vec{S}_2 - 17\vec{S}_1 \cdot \vec{n} \vec{S}_2 \cdot \vec{n} \right) - \frac{G_N}{r^3} \left(\vec{S}_1 \cdot \vec{S}_2 - 3\vec{S}_1 \cdot \vec{n} \vec{S}_2 \cdot \vec{n} \right), \quad (4)
\end{aligned}$$

where we also added last, the LO spin(1)-spin(2) potential at 2PN order.

We can now show that the canonical transformation generated by ($b = c = 1/2$ in [5])

$$g = \frac{G_N}{2r^2} \left[\frac{1}{m_2} \vec{S}_1 \cdot \vec{p}_2 \vec{S}_2 \cdot \vec{n} - \frac{1}{m_1} \vec{S}_2 \cdot \vec{p}_1 \vec{S}_1 \cdot \vec{n} \right] + \frac{G_N}{2r^2} \left[\frac{1}{m_1} \vec{S}_1 \cdot \vec{S}_2 \vec{p}_1 \cdot \vec{n} - \frac{1}{m_2} \vec{S}_2 \cdot \vec{S}_1 \vec{p}_2 \cdot \vec{n} \right], \quad (5)$$

leads to $\delta H_{SS}^{NLO} = 0$ at each order in G_N . Thus the results in [5] do indeed reproduce those in [1]. In a forthcoming paper we will present full details of the spin-spin calculation.

-
- [1] R. A. Porto and I. Z. Rothstein, Phys. Rev. Lett. **97**, 021101 (2006)
 - [2] W. Goldberger and I.Z. Rothstein, Phys. Rev. D **73**, 104029 (2006).
 - [3] W. D. Goldberger and I. Z. Rothstein, Phys. Rev. D **73**, 104030 (2006) [arXiv:hep-th/0511133]. “Les Houches lectures on effective field theories and gravitational radiation,” arXiv:hep-ph/0701129.
 - [4] R. A. Porto, Phys. Rev. D **73**, 104031 (2006)
 - [5] J. Steinhoff, S. Hergt and G. Schafer, arXiv:0712.1716v1 [gr-qc].
 - [6] R. A. Porto, arXiv:gr-qc/0701106.
 - [7] A. Hanson and T. Regge, Ann. Phys. (N.Y.) **87**, 498 (1974).